

Parameterised Subgraph Counting Problems

Kitty Meeks

University of Glasgow

Bergen, 13th May 2015

Joint work with Mark Jerrum (QMUL)

What is a counting problem?

Decision problems

Given a graph G , does G contain a Hamilton cycle?

Given a bipartite graph G , does G contain a perfect matching?

What is a counting problem?

Decision problems

Given a graph G , does G contain a Hamilton cycle?

Given a bipartite graph G , does G contain a perfect matching?

Counting problems

How many Hamilton cycles are there in the graph G ?

How many perfect matchings are there in the bipartite graph G ?

What is a parameterised counting problem?

- Introduced by Flum and Grohe (2004)
- Measure running time in terms of a **parameter** as well as the total input size
- Examples:

What is a parameterised counting problem?

- Introduced by Flum and Grohe (2004)
- Measure running time in terms of a **parameter** as well as the total input size
- Examples:
 - How many vertex-covers of size k are there in G ?

What is a parameterised counting problem?

- Introduced by Flum and Grohe (2004)
- Measure running time in terms of a **parameter** as well as the total input size
- Examples:
 - How many vertex-covers of size k are there in G ?
 - How many k -cliques are there in G ?

What is a parameterised counting problem?

- Introduced by Flum and Grohe (2004)
- Measure running time in terms of a **parameter** as well as the total input size
- Examples:
 - How many vertex-covers of size k are there in G ?
 - How many k -cliques are there in G ?
 - Given a graph G of treewidth at most k , how many Hamilton cycles are there in G ?

The theory of parameterised counting

Efficient algorithms: Fixed parameter tractable (FPT)

Running time $f(k) \cdot n^{O(1)}$

The theory of parameterised counting

Efficient algorithms: Fixed parameter tractable (FPT)

Running time $f(k) \cdot n^{O(1)}$

Intractable problems: #W[1]-hard

A #W[1]-complete problem: **p-#CLIQUE**.

#W[1]-completeness

- To show the problem Π' (with parameter κ') is #W[1]-hard, we give a reduction from a problem Π (with parameter κ) to Π' .

#W[1]-completeness

- To show the problem Π' (with parameter κ') is #W[1]-hard, we give a reduction from a problem Π (with parameter κ) to Π' .
- An fpt Turing reduction from (Π, κ) to (Π', κ') is an algorithm A with an oracle to Π' such that
 - 1 A computes Π ,
 - 2 A is an fpt-algorithm with respect to κ , and
 - 3 there is a computable function $g : \mathbb{N} \rightarrow \mathbb{N}$ such that for all oracle queries “ $\Pi'(y) = ?$ ” posed by A on input x we have $\kappa'(y) \leq g(\kappa(x))$.

In this case we write $(\Pi, \kappa) \leq_T^{fpt} (\Pi', \kappa')$.

Subgraph Counting Model

Let Φ be a family (ϕ_1, ϕ_2, \dots) of functions, such that ϕ_k is a mapping from labelled graphs on k -vertices to $\{0, 1\}$.

Subgraph Counting Model

Let Φ be a family (ϕ_1, ϕ_2, \dots) of functions, such that ϕ_k is a mapping from labelled graphs on k -vertices to $\{0, 1\}$.

p-#INDUCED SUBGRAPH WITH PROPERTY(Φ) (ISWP(Φ))

Input: A graph $G = (V, E)$ and an integer k .

Parameter: k .

Question: What is the cardinality of the set

$$\{(v_1, \dots, v_k) \in V^k : v_1, \dots, v_k \text{ all distinct,} \\ \text{and } \phi_k(G[v_1, \dots, v_k]) = 1\}?$$

- $\mathbf{p\text{-}\#\text{SUB}(\mathcal{H})}$

e.g. $\mathbf{p\text{-}\#\text{CLIQUE}}$, $\mathbf{p\text{-}\#\text{PATH}}$, $\mathbf{p\text{-}\#\text{CYCLE}}$, $\mathbf{p\text{-}\#\text{MATCHING}}$

- $\mathbf{p\text{-}\#SUB(\mathcal{H})}$

e.g. $\mathbf{p\text{-}\#CLIQUE}$, $\mathbf{p\text{-}\#PATH}$, $\mathbf{p\text{-}\#CYCLE}$, $\mathbf{p\text{-}\#MATCHING}$

- $\mathbf{p\text{-}\#CONNECTED INDUCED SUBGRAPH}$

Examples

- $\mathbf{p\text{-}\#SUB(\mathcal{H})}$

e.g. $\mathbf{p\text{-}\#CLIQUE}$, $\mathbf{p\text{-}\#PATH}$, $\mathbf{p\text{-}\#CYCLE}$, $\mathbf{p\text{-}\#MATCHING}$

- $\mathbf{p\text{-}\#CONNECTED INDUCED SUBGRAPH}$

- $\mathbf{p\text{-}\#PLANAR INDUCED SUBGRAPH}$

Examples

- $\mathbf{p\text{-}\#SUB(\mathcal{H})}$
e.g. $\mathbf{p\text{-}\#CLIQUE}$, $\mathbf{p\text{-}\#PATH}$, $\mathbf{p\text{-}\#CYCLE}$, $\mathbf{p\text{-}\#MATCHING}$
- $\mathbf{p\text{-}\#CONNECTED INDUCED SUBGRAPH}$
- $\mathbf{p\text{-}\#PLANAR INDUCED SUBGRAPH}$
- $\mathbf{p\text{-}\#EVEN INDUCED SUBGRAPH}$
 $\mathbf{p\text{-}\#ODD INDUCED SUBGRAPH}$

Complexity Questions

- Is the corresponding decision problem in FPT?
- Is there a fixed parameter algorithm for $\mathbf{p}\text{-}\#\text{INDUCED SUBGRAPH WITH PROPERTY}(\Phi)$?
- Can we approximate $\mathbf{p}\text{-}\#\text{INDUCED SUBGRAPH WITH PROPERTY}(\Phi)$ efficiently?

Approximation Algorithms

An FPTRAS for a parameterised counting problem Π with parameter k is a randomised approximation scheme that takes an instance I of Π (with $|I| = n$), and numbers $\epsilon > 0$ and $0 < \delta < 1$, and in time $f(k) \cdot g(n, 1/\epsilon, \log(1/\delta))$ (where f is any function, and g is a polynomial in n , $1/\epsilon$ and $\log(1/\delta)$) outputs a rational number z such that

$$\mathbb{P}[(1 - \epsilon)\Pi(I) \leq z \leq (1 + \epsilon)\Pi(I)] \geq 1 - \delta.$$

The Diagram of Everything


DECISION

APPROX
COUNTING



EXACT
COUNTING




The Diagram of Everything




The Diagram of Everything



The Diagram of Everything



The Diagram of Everything



Monotone properties I: $\mathbf{p}\text{-}\#\text{SUB}(\mathcal{H})$

Theorem (Arvind & Raman, 2002)

There is an FPTAS for $\mathbf{p}\text{-}\#\text{SUB}(\mathcal{H})$ whenever all graphs in \mathcal{H} have bounded treewidth.

Monotone properties I: $\mathbf{p}\text{-}\#\text{SUB}(\mathcal{H})$

Theorem (Arvind & Raman, 2002)

There is an FPTRAS for $\mathbf{p}\text{-}\#\text{SUB}(\mathcal{H})$ whenever all graphs in \mathcal{H} have bounded treewidth.

Theorem (Curticapean & Marx, 2014)

$\mathbf{p}\text{-}\#\text{SUB}(\mathcal{H})$ is in FPT if all graphs in \mathcal{H} have bounded vertex-cover number; otherwise $\mathbf{p}\text{-}\#\text{SUB}(\mathcal{H})$ is $\#W[1]$ -complete.

Monotone properties II: properties with more than one minimal element

Theorem (Jerrum & M.)

Let Φ be a monotone property, and suppose that there exists a constant t such that, for every $k \in \mathbb{N}$, all minimal graphs satisfying ϕ_k have treewidth at most t . Then there is an FPTRAS for $\mathbf{p}\text{-}\#\text{INDUCED SUBGRAPH WITH PROPERTY}(\Phi)$.

Monotone properties II: properties with more than one minimal element

Theorem (M.)

Suppose that there is no constant t such that, for every $k \in \mathbb{N}$, all minimal graphs satisfying ϕ_k have treewidth at most t . Then $\mathbf{p}\text{-}\#\text{INDUCED SUBGRAPH WITH PROPERTY}(\Phi)$ is $\#W[1]$ -complete.

Monotone properties II: properties with more than one minimal element

Theorem (M.)

Suppose that there is no constant t such that, for every $k \in \mathbb{N}$, all minimal graphs satisfying ϕ_k have treewidth at most t . Then \mathbf{p} -#INDUCED SUBGRAPH WITH PROPERTY(Φ) is #W[1]-complete.

Theorem (Jerrum & M.)

\mathbf{p} -#CONNECTED INDUCED SUBGRAPH is #W[1]-complete.

Non-monotone properties

Theorem

Let Φ be a family (ϕ_1, ϕ_2, \dots) of functions ϕ_k from labelled k -vertex graphs to $\{0, 1\}$ that are not identically zero, such that the function mapping $k \mapsto \phi_k$ is computable. Suppose that

$$|\{|E(H)| : |V(H)| = k \text{ and } \Phi \text{ is true for } H\}| = o(k^2).$$

Then **p-#INDUCED SUBGRAPH WITH PROPERTY(Φ)** is **#W[1]-complete**.

Non-monotone properties

Theorem

Let Φ be a family (ϕ_1, ϕ_2, \dots) of functions ϕ_k from labelled k -vertex graphs to $\{0, 1\}$ that are not identically zero, such that the function mapping $k \mapsto \phi_k$ is computable. Suppose that

$$|\{ |E(H)| : |V(H)| = k \text{ and } \Phi \text{ is true for } H \}| = o(k^2).$$

Then **p-#INDUCED SUBGRAPH WITH PROPERTY(Φ)** is **#W[1]-complete**.

E.g. **p-#PLANAR INDUCED SUBGRAPH**, **p-#REGULAR INDUCED SUBGRAPH**

Even induced subgraphs: FPT???

Theorem (Goldberg, Grohe, Jerrum & Thurley (2010); Lidl & Niederreiter (1983))


Given a graph G , there is a polynomial-time algorithm which computes the number of induced subgraphs of G having an even number of edges.

Even induced subgraphs: decision


Let G be a graph on $n \geq 2^{2k}$ vertices. Then:

- If $k \equiv 0 \pmod{4}$ or $k \equiv 1 \pmod{4}$ then G contains a k -vertex subgraph with an even number of edges.
- If $k \equiv 2 \pmod{4}$ then G contains a k -vertex subgraph with an even number of edges *unless* G is a clique.
- If $k \equiv 3 \pmod{4}$ then G contains a k -vertex subgraph with an even number of edges *unless* G is either a clique or the disjoint union of two cliques.


Even induced subgraphs: exact counting is $\#W[1]$ -complete




Even induced subgraphs: exact counting is $\#W[1]$ -complete



Even induced subgraphs: exact counting is $\#W[1]$ -complete



Even induced subgraphs: exact counting is #W[1]-complete



underlying structure

mask		0	1	1	1	0	0	0	1
		1	1	0	0	0	0	1	1
		1	0	1	0	0	1	0	1
		1	0	0	1	1	0	0	1
		0	0	0	1	0	1	1	1
		0	0	1	0	1	0	1	1
		0	1	0	0	1	1	0	1
		1	1	1	1	1	1	1	1

·


	#
	#
	#
	#
	#
	#
	#
	#
	#

=

·
·
·
·
·
·
·
·
·

University of Glasgow


Even induced subgraphs: exact counting is $\#W[1]$ -complete



underlying structure

$$\text{mask} \begin{pmatrix} \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \end{pmatrix} \cdot \begin{pmatrix} \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ 0 & 1 & 1 & 1 & 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{pmatrix} \cdot \begin{pmatrix} \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \end{pmatrix} = \begin{pmatrix} 4 \\ \cdot \\ \cdot \\ \cdot \\ \cdot \\ \cdot \\ \cdot \\ \cdot \\ \cdot \end{pmatrix}$$


Even induced subgraphs: exact counting is $\#W[1]$ -complete



underlying structure

$$\text{mask} \begin{pmatrix} \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \end{pmatrix} \begin{pmatrix} \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \end{pmatrix} \begin{pmatrix} \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \\ \cdot \\ \cdot \\ \cdot \\ \cdot \\ \cdot \\ \cdot \\ \cdot \end{pmatrix}$$


Even induced subgraphs: exact counting is $\#W[1]$ -complete



underlying structure

$$\text{mask} \begin{pmatrix} \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \end{pmatrix} \begin{pmatrix} \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \end{pmatrix} \begin{pmatrix} \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \\ \triangle \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \\ 2 \\ \cdot \\ \cdot \\ \cdot \\ \cdot \\ \cdot \\ \cdot \\ \cdot \end{pmatrix}$$


Even induced subgraphs: exact counting is $\#W[1]$ -complete



underlying structure

$$\text{mask} \begin{pmatrix} \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \\ \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle & \triangle \end{pmatrix} \cdot \begin{pmatrix} \# \triangle \\ \# \triangle \\ \# \triangle \\ \# \triangle \\ \# \triangle \\ \# \triangle \\ \# \triangle \\ \# \triangle \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \\ 2 \\ 6 \\ \cdot \\ \cdot \\ \cdot \end{pmatrix}$$


Even induced subgraphs: exact counting is #W[1]-complete



underlying structure

$$\text{mask} \begin{pmatrix} \begin{matrix} \text{triangle} \\ \text{V-shape} \\ \text{V-shape} \\ \text{V-shape} \\ \text{V-shape} \\ \text{V-shape} \\ \text{V-shape} \\ \text{V-shape} \\ \text{V-shape} \\ \text{V-shape} \\ \text{V-shape} \end{matrix} & \begin{pmatrix} 0 & 1 & 1 & 1 & 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{pmatrix} \cdot \begin{pmatrix} \# \text{triangle} \\ \# \text{V-shape} \\ \# \text{V-shape} \\ \# \text{V-shape} \\ \# \text{V-shape} \\ \# \text{V-shape} \\ \# \text{V-shape} \\ \# \text{V-shape} \\ \# \text{V-shape} \\ \# \text{V-shape} \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \\ 2 \\ 6 \\ 4 \\ \cdot \\ \cdot \end{pmatrix}
 \end{pmatrix}$$


Even induced subgraphs: exact counting is $\#W[1]$ -complete



underlying structure

{	mask									}	·	()	=	(4)	
		0	1	1	1	0	0	0	1								#		4
		1	1	0	0	0	0	1	1								#		2
		1	0	1	0	0	1	0	1								#		6
		1	0	0	1	1	0	0	1								#		4
		0	0	0	1	0	1	1	1								#		2
		0	0	1	0	1	0	1	1								#		.
		0	1	0	0	1	1	0	1								#		
	1	1	1	1	1	1	1	1		#									

Even induced subgraphs: exact counting is $\#W[1]$ -complete



underlying structure

{	mask								
		0	1	1	1	0	0	0	1
		1	1	0	0	0	0	1	1
		1	0	1	0	0	1	0	1
		1	0	0	1	1	0	0	1
		0	0	0	1	0	1	1	1
		0	0	1	0	1	0	1	1
		0	1	0	0	1	1	0	1
	1	1	1	1	1	1	1	1	

 $\cdot \begin{pmatrix} \# \text{triangle with red top and yellow bottom-right} \\ \# \text{triangle with red top and blue bottom-right} \\ \# \text{triangle with red top and yellow bottom-left} \\ \# \text{triangle with red top and blue bottom-left} \\ \# \text{triangle with red top and yellow bottom-left and blue bottom-right} \\ \# \text{triangle with red top and blue bottom-left and yellow bottom-right} \\ \# \text{triangle with red top} \\ \# \text{triangle with red top and yellow bottom-left and blue bottom-right} \\ \# \text{triangle with red top and blue bottom-left and yellow bottom-right} \\ \# \text{triangle with red top} \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \\ 2 \\ 6 \\ 4 \\ 2 \\ 2 \\ 2 \end{pmatrix}$

Even induced subgraphs: an FPTRAS

Lemma

Suppose that, for each k and any graph G on n vertices, the number of k -vertex (labelled) subgraphs of G that satisfy ϕ_k is either

- 1 zero, or
- 2 at least

$$\frac{1}{g(k)p(n)} \binom{n}{k},$$

where p is a polynomial and g is a computable function.

Then there exists an FPTRAS for $\mathbf{p}\text{-}\#\text{ISWP}(\Phi)$.

Even induced subgraphs: an FPTRAS

Theorem

Let $k \geq 3$ and let G be a graph on $n \geq 2^{2k}$ vertices. Then either G contains no even k -vertex subgraph or else G contains at least

$$\frac{1}{2^{2k^2} k^2 n^2} \binom{n}{k}$$

even k -vertex subgraphs.

Even induced subgraphs: an FPTRAS

Theorem (Erdős and Szekeres)

Let $k \in \mathbb{N}$. Then there exists $R(k) < 2^{2^k}$ such that any graph on $n \geq R(k)$ vertices contains either a clique or independent set on k vertices.

Corollary

Let $G = (V, E)$ be an n -vertex graph, where $n \geq 2^{2^k}$. Then the number of k -vertex subsets $U \subset V$ such that U induces either a clique or independent set in G is at least

$$\frac{(2^{2^k} - k)!}{(2^{2^k})!} \frac{n!}{(n - k)!}.$$

Corollary

Let $G = (V, E)$ be an n -vertex graph, where $n \geq 2^{2^k}$. Then the number of k -vertex subsets $U \subset V$ such that U induces either a clique or independent set in G is at least

$$\frac{(2^{2^k} - k)!}{(2^{2^k})!} \frac{n!}{(n-k)!}.$$

- If at least half of these “interesting” subsets are independent sets, we are done.
- Thus we may assume from now on that G contains at least $\frac{(2^{2^k} - k)!}{2(2^{2^k})!} \frac{n!}{(n-k)!}$ k -cliques.


Even induced subgraphs: an FPTRAS

Definition


Let $A \subset \{1, \dots, k\}$. We say that a k -clique H in G is A -extendible if there are subsets $U \subset V(H)$ and $W \subset V(G) \setminus V(H)$, with $|U| = |W| \in A$, such that $G[(H \setminus U) \cup W]$ has an even number of edges.

- If every k -clique in G is $\{1, 2\}$ -extendible, we are done.
- Thus we may assume from now on that there is at least one k -clique H in G that is not $\{1, 2\}$ -extendible.


Even induced subgraphs: an FPTRAS



Even induced subgraphs: an FPTRAS





Even induced subgraphs: an FPTRAS




Even induced subgraphs: an FPTRAS

If $\binom{k}{2}$ is odd, the following have an even number of edges:




Even induced subgraphs: an FPTRAS

If $\binom{k}{2}$ is odd, the following have an even number of edges:




If $k \equiv 2 \pmod{4}$, this also has an even number of edges:




Even induced subgraphs: an FPTRAS




Even induced subgraphs: an FPTRAS




Even induced subgraphs: an FPTRAS




Even induced subgraphs: an FPTRAS




Even induced subgraphs: an FPTRAS



Even induced subgraphs: an FPTRAS



Even induced subgraphs: an FPTRAS



Open problems

Open problems

- Can similar results be obtained for properties that only hold for graphs H where

$$e(H) \equiv r \pmod{p},$$

for $p > 2$?

Open problems

- Can similar results be obtained for properties that only hold for graphs H where

$$e(H) \equiv r \pmod{p},$$

for $p > 2$?

- What if we consider an arbitrary property that depends only on the number of edges?

THANK YOU