

# Stable Marriage with Ties and Bounded Length Preference Lists

Robert W. Irving, David F. Manlove<sup>\*,†</sup> and Gregg O'Malley<sup>\*</sup>

*Department of Computing Science, University of Glasgow, Glasgow G12 8QQ, UK.*

*Email: {rwi,davidm,gregg}@dcs.gla.ac.uk.*

**Abstract.** We consider variants of the classical stable marriage problem in which preference lists may contain ties, and may be of bounded length. Such restrictions arise naturally in practical applications, such as centralised matching schemes that assign graduating medical students to their first hospital posts. In such a setting, weak stability is the most common solution concept, and it is known that weakly stable matchings can have different sizes. This motivates the problem of finding a maximum cardinality weakly stable matching, which is known to be NP-hard in general. We show that this problem is solvable in polynomial time if each man's list is of length at most 2 (even for women's lists that are of unbounded length). However if each man's list is of length at most 3, we show that the problem becomes NP-hard and not approximable within some  $\delta > 1$ , even if each woman's list is of length at most 4.

**Keywords:** Stable marriage problem; ties; incomplete lists; NP-hardness; polynomial-time algorithm

## 1 Introduction

The Stable Marriage problem (SM) was introduced in the seminal paper of Gale and Shapley [3]. In its classical form, an instance of SM involves  $n$  men and  $n$  women, each of whom specifies a *preference list*, which is a total order on the members of the opposite sex. A *matching*  $M$  is a set of (man,woman) pairs such that each person belongs to exactly one pair. If  $(m, w) \in M$ , we say that  $w$  is  $m$ 's *partner* in  $M$ , and vice versa, and we write  $M(m) = w$ ,  $M(w) = m$ .

We say that a person  $x$  *prefers*  $y$  to  $y'$  if  $y$  precedes  $y'$  on  $x$ 's preference list. A matching  $M$  is *stable* if it admits no *blocking pair*, namely a pair  $(m, w)$  such that  $m$  prefers  $w$  to  $M(m)$  and  $w$  prefers  $m$  to  $M(w)$ . Gale and Shapley [3] proved that every instance of SM admits at least one stable matching, and described an algorithm – the Gale / Shapley algorithm – that finds such a matching in time that

---

<sup>\*</sup>Supported by Engineering and Physical Sciences Research Council grant GR/R84597/01.

<sup>†</sup>Supported by Royal Society of Edinburgh/Scottish Executive Personal Research Fellowship.

is linear in the input size. In general, there may be many stable matchings (in fact exponentially many in  $n$ ) for a given instance of SM [13].

**Incomplete lists.** A variety of extensions of the basic problem have been studied. In the Stable Marriage problem with Incomplete lists (SMI), the numbers of men and women need not be the same, and each person's preference list consists of a subset of the members of the opposite sex in strict order. A pair  $(m, w)$  is *acceptable* if each member of the pair appears on the preference list of the other. A *matching*  $M$  is now a set of acceptable pairs such that each person belongs to at most one pair. In this context,  $(m, w)$  is a blocking pair for a matching  $M$  if (a)  $(m, w)$  is an acceptable pair, (b)  $m$  is either unmatched or prefers  $w$  to  $M(m)$ , and likewise (c)  $w$  is either unmatched or prefers  $m$  to  $M(w)$ . As in the classical case, there is always at least one stable matching for an instance of SMI, and it is straightforward to extend the Gale / Shapley algorithm to give a linear-time algorithm for this case. Again, there may be many different stable matchings, but Gale and Sotomayor [4] showed that every stable matching for a given SMI instance has the same size and matches exactly the same set of people.

**Ties.** An alternative extension of SM arises if preference lists are allowed to contain *ties*. In the Stable Marriage problem with Ties (SMT) each person's preference list is a partial order over the members of the opposite sex in which indifference is transitive. In other words, each person  $p$ 's list can be viewed as a sequence of ties, each of length  $\geq 1$ ;  $p$  prefers each member of a tie to everyone in any subsequent tie, but is indifferent between the members of any single tie. In this context, three definitions of stability have been proposed [6, 11]. Among these three stability criteria, it is *weak stability* that has received the most attention in the literature [15, 18, 8, 7, 9, 10, 16, 17]. A matching  $M$  is *weakly stable* if there is no pair  $(m, w)$ , each of whom prefers the other to his/her partner in  $M$ . For a given instance of SMT, a weakly stable matching is bound to exist, and can be found in linear time by breaking all ties in an arbitrary way (i.e. by strictly ranking the members of each tie arbitrarily) and applying the Gale / Shapley algorithm.

**Ties and incomplete lists.** If we allow both of the above extensions of the classical problem simultaneously, we obtain the Stable Marriage problem with Ties and Incomplete lists (SMTI). In this context a matching  $M$  is weakly stable if there is no acceptable pair  $(m, w)$ , each of whom is either unmatched in  $M$  or prefers the other to his/her partner in  $M$ . Once again, it is easy to find a weakly stable matching, merely by breaking all the ties in an arbitrary way and applying the Gale / Shapley algorithm. However, the ways in which ties are broken will, in general, affect the size of the resulting matching. It is therefore natural to consider MAX SMTI, the problem of finding a maximum cardinality weakly stable matching (henceforth a maximum weakly stable matching), given an instance of SMTI. MAX SMTI turns out to be NP-hard, even under quite severe restrictions on the number and lengths of ties [18]. Specifically, NP-hardness holds even if ties occur in the men's preference lists only, each tie is of length 2, and each tie comprises the whole of the list in which it appears [18]. (Note that, in the SMTI instance constructed by the reduction in [18], there are men with strictly ordered preference lists of length at least 3.)

**The Hospitals/Residents problem.** The Hospitals/Residents problem (HR) is a many-to-one generalisation of SMI, so called because of its application in centralised matching schemes for the allocation of graduating medical students, or residents, to hospitals [20]. The best known such scheme is the National Resident Matching Program (NRMP) [19] in the US, but similar schemes exist in Canada [1], in Scotland [12, 21], and in a variety of other countries and contexts. In fact, this extension of SM was also discussed by Gale and Shapley under the name of the College Admissions problem [3]. In an instance of HR, each resident has a preference list containing a subset of the hospitals, and each hospital ranks the residents for which it is acceptable. In addition, each hospital has a *quota* of available posts. In this context, a matching is a set of acceptable (resident,hospital) pairs so that each resident appears in at most one pair and each hospital in a number of pairs that is bounded by its quota. The definition of stability is easily extended to this more general setting (see [6] for details). It is again the case that every problem instance admits at least one stable matching [3], and that all stable matchings have the same size [4]. Clearly SMI is equivalent to the special case of HR in which each hospital has a quota of 1.

The Hospitals / Residents problem with Ties (HRT) allows arbitrary ties in the preference lists. The definition of weak stability can be extended in a natural way to the HRT context [14]. Since HRT is clearly an extension of SMTI, the hardness results for weak stability problems in the latter extend to the former. These results have potentially important implications for large-scale real-world matching schemes. It is unreasonable to expect, say, a large hospital to rank in strict order all of its many applicants, and any artificial rankings, whether submitted by the hospitals themselves, or imposed by the matching scheme administrators, may have significant implications for the number of residents assigned in a stable matching.

**Bounded length preference lists.** In the context of many large-scale matching schemes, the preference lists of at least one set of agents tend to be short. For example, until recently, students participating in the Scottish medical matching scheme [12, 21] were required to rank just three hospitals in order of preference. This naturally leads to the question of whether the problem of finding a maximum weakly stable matching becomes simpler when preference lists on one or both sides have bounded length.

Let  $(p, q)$ -MAX SMTI denote the restriction of MAX SMTI in which each man's list is of length at most  $p$  and each woman's list is of length at most  $q$ . We use  $p = \infty$  or  $q = \infty$  to denote the possibility that the men's lists or women's lists are of unbounded length, respectively. Halldorsson et al. [7] showed that  $(4, 7)$ -MAX SMTI is NP-hard and not approximable within some  $\delta > 1$  unless  $P=NP$ . Halldorsson et al. [9] gave an alternative reduction from Minimum Vertex Cover to MAX SMTI, showing that the latter problem is not approximable within  $\frac{21}{19}$  unless  $P=NP$ . By starting from the NP-hard restriction of Minimum Vertex Cover to graphs of maximum degree 3 [5], the same reduction shows NP-hardness for  $(5, 5)$ -MAX SMTI.

In this paper we consider other values of  $p$  and  $q$ , to narrow down the search for the 'borderline' between polynomial-time solvability and NP-hardness for  $(p, q)$ -MAX SMTI. We show in Section 2 that  $(2, \infty)$ -MAX SMTI is polynomial-time solvable using a combination of an adapted version of the Gale / Shapley algorithm together with a reduction to the Assignment problem. By contrast, in Section 3 we show

that (3,4)-MAX SMTI is NP-hard and not approximable within some  $\delta > 1$  unless  $P=NP$ . Finally, in Section 4 we present some concluding remarks.

## 2 Algorithm for $(2, \infty)$ -MAX SMTI

In this section we present a polynomial-time algorithm for MAX SMTI where the preference lists of both men and women may contain ties, the men’s lists are of length at most 2 and the women’s lists are of unbounded length. Let  $I$  be an instance of this problem, where  $n_1$  and  $n_2$  are the numbers of men and women in  $I$  respectively.

Consider the algorithm  $(2, \infty)$ -MAX-SMTI-**alg** shown in Figure 1. The algorithm consists of three phases, where each phase is highlighted in the figure. We use the term *reduced lists* to refer to participants’ lists after any deletions made by the algorithm. Phase 1 of  $(2, \infty)$ -MAX-SMTI-**alg** is a simple extension of the Gale / Shapley algorithm, and is used to delete certain (man,woman) pairs that can never be part of any weakly stable matching. To “delete the pair  $(m_i, w_j)$ ”, we delete  $m_i$  from  $w_j$ ’s list and delete  $w_j$  from  $m_i$ ’s list. Phase 1 proceeds as follows. All men are initially unmarked. While some man  $m_i$  remains unmarked and  $m_i$  has a non-empty reduced list, we set  $m_i$  to be marked – it is possible that  $m_i$  may again become unmarked at a later stage of the execution. If  $m_i$ ’s reduced list is not a tie of length 2, we let  $w_j$  be the woman in first position in  $m_i$ ’s reduced list. Then, for each strict successor  $m_k$  of  $m_i$  on  $w_j$ ’s list, we delete the pair  $(m_k, w_j)$  and set  $m_k$  to be unmarked (regardless of whether or not he was already marked).

We remark that the following situation may occur during phase 1. Suppose that some man  $m_i$  is indifferent between two women  $w_j$  and  $w_k$  on his original preference list, and suppose that during some iteration of the while loop he becomes marked. We note that the algorithm does not delete the strict successors of  $m_i$  on  $w_j$ ’s list at this stage. Now suppose that, during a subsequent loop iteration, the pair  $(m_i, w_k)$  is deleted. Then  $m_i$  becomes unmarked, only to be re-marked during a subsequent loop iteration. This re-marking results in the deletions of all pairs  $(m_r, w_j)$ , where  $m_r$  is a strict successor of  $m_i$  on  $w_j$ ’s list, as required.

In phase 2 we construct a weighted bipartite graph  $G$  and find a minimum cost maximum matching in  $G$  using the algorithm in [2]. The graph  $G$  is constructed using Algorithm **BuildGraph** shown in Figure 2. That is, each man and woman is represented by a vertex in  $G$ , and for each man  $m_i$  on woman  $w_j$ ’s reduced list, we add an edge from  $m_i$  to  $w_j$  with cost  $rank(w_j, m_i)$ , where  $rank(w_j, m_i)$  is the rank of  $m_i$  on  $w_j$ ’s reduced list (i.e. 1 plus the number of strict predecessors of  $m_i$  on  $w_j$ ’s reduced list). We then find a minimum cost maximum matching  $M_G$  in  $G$ .

In general, after phase 2,  $M_G$  need not be weakly stable in  $I$ . In particular, some man  $m_i$  who has a reduced list of length 2 that is strictly ordered may be assigned to his second-choice woman  $w_k$  in  $M_G$ , while his first-choice woman  $w_j$  may be unassigned in  $M_G$ . Clearly  $(m_i, w_j)$  blocks such a matching. To obtain a weakly stable matching  $M$  from  $M_G$  we execute phase 3. Initially  $M$  is set to be equal to  $M_G$ . Next, we move each such  $m_i$  to his first-choice woman. We note that  $m_i$  must be in the *tail* of  $w_j$ ’s reduced list (this is the set of one or more entries tied in last place on  $w_j$ ’s reduced list) since  $m_i$  must have been marked during Phase 1, causing all strict successors of  $m_i$  on  $w_j$ ’s list to be deleted. Further, we note

```

/* Phase 1 */
set all men to be unmarked;
while (some man  $m_i$  is unmarked and
 $m_i$  has a non-empty reduced list) do
    set  $m_i$  to be marked;
    if  $m_i$ 's reduced list is not a tie of length 2 then
         $w_j :=$  woman in first position on  $m_i$ 's reduced list;
        for each strict successor  $m_k$  of  $m_i$  on  $w_j$ 's list do
            set  $m_k$  to be unmarked;
            delete the pair  $(m_k, w_j)$ ;

/* Phase 2 */
 $G :=$  BuildGraph();
 $M_G :=$  minimum cost maximum matching in  $G$ ;

/* Phase 3 */
 $M := M_G$ ;
while (there exists a man  $m_i$  who is assigned
    to his second-choice woman  $w_k$  in  $M$ 
    and his first-choice woman  $w_j$  is unassigned in  $M$ ) do
     $M := M \setminus \{(m_i, w_k)\}$ ;
     $M := M \cup \{(m_i, w_j)\}$ ;
return  $M$ ;

```

Figure 1: Algorithm  $(2, \infty)$ -MAX-SMTI-**alg**.

that there may exist more than one such man in  $w_j$ 's tail who satisfies the above criterion. Moreover when  $m_i$  moves to  $w_j$ ,  $w_k$  becomes unassigned in  $M$ . As a result, there may be some other man  $m_r$  (who strictly ranks  $w_k$  in first place) who now satisfies the loop condition. This process is repeated until no such man exists. Upon termination of phase 3 we will show that the matching  $M$  returned is a maximum weakly stable matching.

We begin by showing that the algorithm  $(2, \infty)$ -MAX-SMTI-**alg** terminates. It is easy to see that each of phases 1 and 2 is bound to terminate. The following lemma shows that the same is true for phase 3.

**Lemma 1.** *Phase 3 of  $(2, \infty)$ -MAX-SMTI-**alg** terminates.*

*Proof.* We show that the while loop terminates during an execution  $E$  of phase 3. For, at a given iteration of the while loop of phase 3, let  $m_i$  be some man assigned to his second-choice woman  $w_k$  in  $M$  and suppose that his first-choice woman  $w_j$  is unassigned in  $M$ , where  $m_i$ 's reduced list is of length 2 and is strictly ordered. Then during  $E$ ,  $m_i$  switches from  $w_k$  to  $w_j$ . Hence each such  $m_i$  must strictly improve (in fact  $m_i$  can only improve at most once). Therefore since the number of men is finite, phase 3 is bound to terminate.  $\square$

We next show that phase 1 of  $(2, \infty)$ -MAX-SMTI-**alg** never deletes a *weakly stable pair*, which is a (man, woman) pair that belongs to some weakly stable matching in  $I$ .

```

 $V := \mathcal{M} \cup \mathcal{W};$ 
 $E := \emptyset;$ 
for each man  $m_i \in \mathcal{M}$  do
    for each woman  $w_j$  on  $m_i$ 's reduced list do
         $E := E \cup \{(m_i, w_j)\};$ 
         $cost(m_i, w_j) := rank(w_j, m_i);$ 
 $G := (V, E);$ 
return  $G;$ 

```

Figure 2: Algorithm BuildGraph.

**Lemma 2.** *The algorithm  $(2, \infty)$ -MAX-SMTI-**alg** never deletes a weakly stable pair.*

*Proof.* Let  $(m_i, w_j)$  be a pair deleted during an execution  $E$  of  $(2, \infty)$ -MAX-SMTI-**alg** such that  $(m_i, w_j) \in M$ , where  $M$  is a weakly stable matching in  $I$ . Without loss of generality suppose this is the first weakly stable pair deleted during  $E$ . Then  $m_i$  was deleted from  $w_j$ 's list during some iteration  $q$  of the while loop of phase 1 during  $E$ . This deletion was made as a result of  $w_j$  being in first position in the reduced list of some man  $m_r$ , where  $m_r$ 's reduced list was not a tie of length 2, and  $w_j$  prefers  $m_r$  to  $m_i$ . Then in  $M$ ,  $m_r$  must obtain a woman  $w_s$  whom he prefers to  $w_j$ , otherwise  $(m_r, w_j)$  blocks  $M$ . Therefore during  $E$ ,  $(m_r, w_s)$  must have already been deleted before iteration  $q$ , a contradiction.  $\square$

Finally we prove that the matching returned by  $(2, \infty)$ -MAX-SMTI-**alg** is weakly stable in  $I$ .

**Lemma 3.** *The matching returned by algorithm  $(2, \infty)$ -MAX-SMTI-**alg** is weakly stable in  $I$ .*

*Proof.* Suppose for a contradiction that the matching  $M$  output by the algorithm  $(2, \infty)$ -MAX-SMTI-**alg** is not weakly stable. Then there exists a pair  $(m_i, w_j)$  that blocks  $M$ . We consider the following four cases corresponding to a blocking pair.

*Case (i):* both  $m_i$  and  $w_j$  are unassigned in  $M$ . Then  $m_i$  is unassigned in  $M_G$ , and either  $w_j$  is unassigned in  $M_G$  or becomes unassigned during phase 3. First suppose that  $w_j$  is unassigned in  $M_G$ . Then the size of the matching  $M_G$  could be increased by adding the edge  $(m_i, w_j)$  to  $M_G$ , contradicting the maximality of  $M_G$ . Now suppose that  $w_j$  became unassigned as a result of phase 3. Let  $m_{p_1}$  denote  $w_j$ 's partner in  $M_G$ . Then during phase 3,  $m_{p_1}$  must have become assigned to his first-choice woman  $w_{q_1}$ . Suppose  $w_{q_1}$  was unassigned in  $M_G$ . Then we can find a larger matching by augmenting along the path  $(m_i, w_j), (w_j, m_{p_1}), (m_{p_1}, w_{q_1})$ , contradicting the maximality of  $M_G$ . Therefore  $w_{q_1}$  must have been assigned in  $M_G$  and became unassigned as a result of phase 3. Hence the man  $m_{p_2}$ , to whom  $w_{q_1}$  was assigned in  $M_G$ , switched to his first-choice woman  $w_{q_2}$ . Using an argument similar to that above for  $w_{q_1}$ , we can show that  $w_{q_2}$  must be assigned in  $M_G$ . Therefore some man switched from  $w_{q_2}$  during phase 3 to his first-choice woman. If we continue this process, since each man must strictly improve and the number of men is finite, there exists a finite number of women that can become unassigned as a result of phase 3. Hence at some point there exists a man  $m_{p_r}$  who switches to his first-choice woman  $w_{q_r}$  and

$w_{q_r}$  was already unassigned in  $M_G$ . We can then construct an augmenting path in  $G$  of the form  $(m_i, w_j), (w_j, m_{p_1}), (m_{p_1}, w_{q_1}), (w_{q_1}, m_{p_2}), (m_{p_2}, w_{q_2}), \dots, (m_{p_r}, w_{q_r})$ , which contradicts the maximality of  $M_G$ .

*Case (ii):*  $m_i$  is unassigned in  $M$  and  $w_j$  prefers  $m_i$  to her assignee  $m_k$  in  $M$ . Then  $m_i$  is unassigned in  $M_G$ . Suppose that  $w_j$  is assigned to  $m_k$  in  $M_G$ . As  $w_j$  prefers  $m_i$  to  $m_k$ , we could obtain a matching with a smaller cost, but with the same size, by removing  $(m_k, w_j)$  and adding  $(m_i, w_j)$  to  $M_G$ , a contradiction. Now suppose that  $w_j$  is not assigned to  $m_k$  in  $M_G$ . Then  $w_j$  is either unassigned in  $M_G$  or  $w_j$  is assigned in  $M_G$  to  $m_r$ , where  $m_r \neq m_k$  and  $m_r \neq m_i$ . If  $w_j$  is unassigned in  $M_G$ , we contradict the maximality of  $M_G$ . Now suppose  $w_j$  is assigned to  $m_r$  in  $M_G$ . Then since  $w_j$  is no longer assigned to  $m_r$  in  $M$ ,  $m_r$  must have switched to his first-choice woman  $w_s$  during phase 3. Therefore either  $w_s$  is unassigned in  $M_G$  or  $w_s$  became unassigned as a result of some man switching from  $w_s$  to his first-choice woman. Again using a similar argument to that in Case (i) we obtain an augmenting path that contradicts the maximality of  $M_G$ .

*Case (iii):*  $m_i$  is assigned to  $w_s$  in  $M$  and  $m_i$  prefers  $w_j$  to  $w_s$  and  $w_j$  is unassigned in  $M$ . Thus clearly  $m_i$ 's list is of length 2 and does not contain a tie, and  $w_j$  is  $m_i$ 's first-choice woman. In this situation the loop condition of phase 3 is satisfied. Therefore since the algorithm terminates (Lemma 1) this situation can never arise.

*Case (iv):*  $m_i$  is assigned to  $w_s$  in  $M$  and  $m_i$  prefers  $w_j$  to  $w_s$ , and  $w_j$  is assigned to  $m_r$  in  $M$  and  $w_j$  prefers  $m_i$  to  $m_r$ . Thus again  $m_i$ 's list cannot contain a tie, and  $w_j$  is his first-choice woman. Therefore either  $m_i$  proposed to  $w_j$  during phase 1 or  $w_j$  was deleted from  $m_i$ 's list. Hence  $m_r$  would have been deleted from  $w_j$ 's list during phase 1, so it is then impossible that  $(m_r, w_j) \in M$ .  $\square$

Since phase 1 of the algorithm never deletes a weakly stable pair (by Lemma 2), a maximum weakly stable matching must consist of (man, woman) pairs that belong to the reduced lists. We next note that  $G$  is constructed from the reduced lists, and since we find a maximum matching in  $G$ , the matching output by the algorithm must indeed be a maximum weakly stable matching (by Lemma 3, and since phase 3 does not change the size of the matching output by the algorithm: every man matched in  $M_G$  is also matched in  $M$ ).

The time complexity of the algorithm is dominated by finding the minimum cost maximum matching in  $G = (V, E)$ . The required matching in  $G$  can be constructed in  $O(\sqrt{|E|}|V| \log |V|)$  time [2]. Let  $n = |V| = n_1 + n_2$ . Since  $|E| \leq 2n_1 = O(n)$ , it follows that  $(2, \infty)$ -MAX-SMTI-**alg** has time complexity  $O(n^{\frac{3}{2}} \log n)$ .

We summarise the results of this section in the following theorem.

**Theorem 4.** *Given an instance  $I$  of  $(2, \infty)$ -MAX SMTI, algorithm  $(2, \infty)$ -MAX-SMTI-**alg** returns a weakly stable matching of maximum size in  $O(n^{\frac{3}{2}} \log n)$  time, where  $n$  is the total number of men and women in  $I$ .*

### 3 NP-hardness of $(3, 4)$ -MAX SMTI

In this section we show that, in contrast to the case for  $(2, \infty)$ -MAX SMTI,  $(3, 4)$ -MAX SMTI is NP-hard and not approximable within  $\delta$ , for some  $\delta > 1$ , unless  $P=NP$ . Our

proof involves a reduction from a problem involving matchings in graphs. A matching  $M$  in a graph  $G$  is said to be *maximal* if no proper superset of  $M$  is a matching in  $G$ . Define MIN-MM to be the problem of finding a minimum cardinality maximal matching, given a graph  $G$ . By [7, Theorem 1], MIN-MM is not approximable within some  $\delta_0 > 1$  unless  $P=NP$ . The result holds even for subdivision graphs of cubic graphs. (Given a graph  $G$ , the *subdivision graph* of  $G$ , denoted by  $S(G)$ , is obtained by subdividing each edge  $\{u, w\}$  of  $G$  in order to obtain two edges  $\{u, v\}$  and  $\{v, w\}$  of  $S(G)$ , where  $v$  is a new vertex.)

**Theorem 5.** *(3, 4)-MAX SMTI is NP-hard and not approximable within  $\delta$ , for some  $\delta > 1$ , unless  $P=NP$ .*

*Proof.* Let  $G$  be an instance of MIN-MM restricted to subdivision graphs of cubic graphs. Then  $G = (U, W, E)$  is a bipartite graph where, without loss of generality, each vertex in  $U$  has degree 2 and each vertex in  $W$  has degree 3. Let  $U = \{m_1, \dots, m_s\}$  and let  $W = \{w_1, \dots, w_t\}$ . For each vertex  $m_i \in U$ , let  $W_i$  denote the two vertices adjacent to  $m_i$  in  $G$ . Similarly for each vertex  $w_j \in W$ , let  $U_j$  denote the three vertices adjacent to  $w_j$  in  $G$ . We construct an instance  $I$  of (3, 4)-MAX SMTI as follows: let  $U \cup X$  be the set of men and let  $W \cup Y$  be the set of women, where  $X = \{x_1, \dots, x_t\}$  and  $Y = \{y_1, \dots, y_s\}$ . The preference lists of the men and women in  $I$  are as follows:

$$\begin{array}{ll} m_i : (W_i) & y_i \quad (1 \leq i \leq s) \\ x_i : w_i & (1 \leq i \leq t) \end{array} \qquad \begin{array}{ll} w_j : (U_j) & x_j \quad (1 \leq j \leq t) \\ y_j : m_j & (1 \leq j \leq s) \end{array}$$

In a given preference list, entries within round brackets are tied. Clearly the length of each man's preference list is at most 3, whilst the length of each woman's preference list is at most 4. We claim that  $s^+(I) = s + t - \beta_1^-(G)$ , where  $s^+(I)$  denotes the maximum size of a weakly stable matching in  $I$  and  $\beta_1^-(G)$  denotes the minimum size of a maximal matching in  $G$ .

For suppose that  $G$  has a maximal matching  $M$ , where  $|M| = \beta_1^-(G)$ . We construct a matching  $M'$  in  $I$  as follows. Initially let  $M' = M$ . There remain  $s - |M|$  men in  $U$  that are unmatched in  $M'$ ; denote these men by  $m_{i_r}$  ( $1 \leq i \leq s - |M|$ ), and add  $(m_{i_r}, y_{i_r})$  to  $M'$  for each such  $m_{i_r}$ . Finally there remain  $t - |M|$  women in  $W$  that are unmatched in  $M'$ ; denote these women by  $w_{j_r}$  ( $1 \leq r \leq t - |M|$ ), and add  $(x_{j_r}, w_{j_r})$  to  $M'$  for each such  $w_{j_r}$ . Clearly  $M'$  is a matching in  $I$  such that  $|M'| = |M| + (s - |M|) + (t - |M|) = s + t - \beta_1^-(G)$ . It is straightforward to verify that  $M'$  is weakly stable in  $I$ , and hence  $s^+(I) \geq s + t - \beta_1^-(G)$ .

Conversely suppose that  $M'$  is a weakly stable matching in  $I$ , where  $|M'| = s^+(I)$ . Let  $M = M' \cap E$ . The weak stability of  $M'$  in  $I$  implies that  $M$  is maximal in  $G$ . Moreover, at most  $t - |M|$  women in  $W$  are matched in  $M'$  to men in  $X$ , and at most  $s - |M|$  men in  $U$  are matched in  $M'$  to women in  $Y$ , and hence  $|M'| \leq |M| + (t - |M|) + (s - |M|) = s + t - |M|$ . Thus  $s^+(I) \leq s + t - \beta_1^-(G)$ . Hence the claim is established.

Theorem 1 of [7] shows that it is NP-hard to distinguish between the cases that  $\beta_1^-(G) \leq c_0 m$  and  $\beta_1^-(G) > \delta_0 c_0 m$ , where  $c_0 > 0$  is some constant and  $m = |E|$ . Now if  $\beta_1^-(G) \leq c_0 m$  then  $s^+(I) \geq cs$ , whilst if  $\beta_1^-(G) > \delta_0 c_0 m$  then  $s^+(I) < \delta cs$ , where  $c = (5 - 6c_0)/2$  and  $\delta = (5 - 6\delta_0 c_0)/(5 - 6c_0)$ . The result follows by Theorem 1 and Proposition 4 of [7].  $\square$



## 4 Concluding remarks

In this paper we have presented a polynomial-time algorithm for  $(2, \infty)$ -MAX SMTI, but have shown that, by contrast,  $(3, 4)$ -MAX SMTI is NP-hard and not approximable within some  $\delta > 1$ . This leaves open the complexity of  $(3, 3)$ -MAX SMTI, the resolution of which would shed further light on the boundary between polynomial-time solvability and NP-hardness for variants of SMTI involving bounded-length preference lists.

Another open problem concerns the complexity of  $(3, \infty)$ -MAX SMTI where ties belong to the preference lists of one sex only. Our reduction that establishes the NP-hardness of  $(3, 4)$ -MAX SMTI introduces ties in the preference lists of both sexes. However in many practical applications (such as assigning medical students to hospitals), if the preference lists of one side (e.g. the residents) are short (as in the SFAS application [21]), then it is reasonable for a matching scheme administrator to insist that these lists should be strictly ordered. This motivates the consideration of  $(3, \infty)$ -MAX SMTI in which ties belong to the women's side only.

For those variants of  $(p, q)$ -MAX SMTI that are NP-hard, it remains to investigate the existence of approximation algorithms for these problems that improve on the performance guarantees of those that have already been formulated for the general SMTI case (with no assumptions on the lengths of the preference lists) [18, 9, 16, 17].

Finally, the natural extension of  $(p, q)$ -MAX SMTI to the many-one HRT case may be formulated: we denote this problem by  $(p, q)$ -MAX HRT. It remains to extend the algorithm for  $(2, \infty)$ -MAX SMTI to the case of  $(2, \infty)$ -MAX HRT or prove that the latter problem is NP-hard. Clearly Theorem 5 implies that  $(3, 4)$ -MAX HRT is also NP-hard and not approximable within some  $\delta > 1$ .

## References

- [1] <http://www.carms.ca> (Canadian Resident Matching Service website).
- [2] H.N. Gabow and R.E. Tarjan. Faster scaling algorithms for network problems. *SIAM Journal on Computing*, 18(5):1013–1036, 1989.
- [3] D. Gale and L.S. Shapley. College admissions and the stability of marriage. *American Mathematical Monthly*, 69:9–15, 1962.
- [4] D. Gale and M. Sotomayor. Some remarks on the stable matching problem. *Discrete Applied Mathematics*, 11:223–232, 1985.
- [5] M.R. Garey, D.S. Johnson, and L. Stockmeyer. Some simplified NP-complete graph problems. *Theoretical Computer Science*, 1:237–267, 1976.
- [6] D. Gusfield and R.W. Irving. *The Stable Marriage Problem: Structure and Algorithms*. MIT Press, 1989.
- [7] M. Halldórsson, R.W. Irving, K. Iwama, D.F. Manlove, S. Miyazaki, Y. Morita, and S. Scott. Approximability results for stable marriage problems with ties. *Theoretical Computer Science*, 306(1-3):431–447, 2003.

- [8] M. Halldórsson, K. Iwama, S. Miyazaki, and Y. Morita. Inapproximability results on stable marriage problems. In *Proceedings of LATIN 2002: the Latin-American Theoretical INformatics symposium*, volume 2286 of *Lecture Notes in Computer Science*, pages 554–568. Springer-Verlag, 2002.
- [9] M.M. Halldórsson, K. Iwama, S. Miyazaki, and H. Yanagisawa. Improved approximation of the stable marriage problem. In *Proceedings of ESA 2003: the 11th Annual European Symposium on Algorithms*, volume 2832 of *Lecture Notes in Computer Science*, pages 266–277. Springer-Verlag, 2003.
- [10] M.M. Halldórsson, K. Iwama, S. Miyazaki, and H. Yanagisawa. Randomized approximation of the stable marriage problem. *Theoretical Computer Science*, 325(3):439–465, 2004.
- [11] R.W. Irving. Stable marriage and indifference. *Discrete Applied Mathematics*, 48:261–272, 1994.
- [12] R.W. Irving. Matching medical students to pairs of hospitals: a new variation on a well-known theme. In *Proceedings of ESA '98: the Sixth European Symposium on Algorithms*, volume 1461 of *Lecture Notes in Computer Science*, pages 381–392. Springer-Verlag, 1998.
- [13] R.W. Irving and P. Leather. The complexity of counting stable marriages. *SIAM Journal on Computing*, 15(3):655–667, 1986.
- [14] R.W. Irving, D.F. Manlove, and S. Scott. The Hospitals/Residents problem with Ties. In *Proceedings of SWAT 2000: the 7th Scandinavian Workshop on Algorithm Theory*, volume 1851 of *Lecture Notes in Computer Science*, pages 259–271. Springer-Verlag, 2000.
- [15] K. Iwama, D. Manlove, S. Miyazaki, and Y. Morita. Stable marriage with incomplete lists and ties. In *Proceedings of ICALP '99: the 26th International Colloquium on Automata, Languages, and Programming*, volume 1644 of *Lecture Notes in Computer Science*, pages 443–452. Springer-Verlag, 1999.
- [16] K. Iwama, S. Miyazaki, and K. Okamoto. A  $(2 - c\frac{\log n}{n})$ -approximation algorithm for the stable marriage problem. In *Proceedings of SWAT 2004: the 9th Scandinavian Workshop on Algorithm Theory*, volume 3111 of *Lecture Notes in Computer Science*, pages 349–361. Springer-Verlag, 2004.
- [17] K. Iwama, S. Miyazaki, and N. Yamauchi. A  $(2 - c\frac{1}{\sqrt{n}})$ -approximation algorithm for the stable marriage problem. In *Proceedings of ISAAC 2005: the 16th Annual International Symposium on Algorithms and Computation*, volume 3827 of *Lecture Notes in Computer Science*, pages 902–914. Springer-Verlag, 2005.
- [18] D.F. Manlove, R.W. Irving, K. Iwama, S. Miyazaki, and Y. Morita. Hard variants of stable marriage. *Theoretical Computer Science*, 276(1-2):261–279, 2002.
- [19] <http://www.nrmp.org/> (National Resident Matching Program website).

- [20] A.E. Roth. The evolution of the labor market for medical interns and residents: a case study in game theory. *Journal of Political Economy*, 92(6):991–1016, 1984.
- [21] <http://www.nes.scot.nhs.uk/sfas/> (Scottish Foundation Allocation Scheme website).